

**SEAN ROBINSON, MD**  
**CAPITAL ORTHOPEDICS**

**Knee Arthroscopy**  
**Postoperative instructions**

- **Weight bearing**
  - You are weight bearing as tolerated after surgery
  - Use crutches at all times until you feel comfortable bearing weight independently.
  - You do **NOT** require a brace
    - You can use a neoprene sleeve if desired
  
- **Movement**
  - It is beneficial to change positions often after knee arthroscopy.
  - **DO NOT STAY IN BED.** We recommend at least 10 minutes of walking every hour.
  
- **Sleep**
  - Keep your leg elevated.
  
- **Ice**
  - Use bagged ice **as much as possible** when you get home at intervals of 30 minutes on and off. You should use ice for approximately two weeks. Do not use ice while you are sleeping.
    - Place the ice onto the knee over a thin layer of clothing or a towel, but never directly on the skin.
  
- **Wound care, Dressings/Bandages**
  - Leave surgical bandage on and do not shower for 48 HOURS
  - After 48 hours, remove bandages and gauze, but leave steri-strips (white tape) in place during showers.
  - Its normal to see bloody soaked fluid on the bandages.
  - After showers, leave the incisions open to air with steri-strips in place and gently pat dry. You may place BAND-AIDS over the incisions
  - Do not apply lotions or ointments to the incisions.
  - Do not soak incisions in any pool/spa/bath water until 3 weeks after postoperatively or until your incisions are completely closed.

**After surgery, call the clinic if you have....**

- A fever higher than 101° (38.3° C).
- Changes in your incision: opening, drainage, redness.
- Numbness or tingling or loss of function of your leg.
  - Numbness around the incisions is normal.
- Increased pain or swelling in your calf.
- Nausea or other side effect not controlled by the medications you are given.

- **CALL 911 IMMEDIATELY IF YOU HAVE CHEST PAIN OR SHORTNESS OF BREATH.**